


NEET - SS

SURGICAL GROUP

HNM NOTES

(High - Yield Notes Module wise)



- HIGH YIELD
- MODULE WISE
- HANDWRITTEN

Presented By:
STEM-S

NEET-SS SURGICAL GROUP

Module-WiseNotes

DETAILED INDEX

GENERAL SURGERY

Injury & Wound Healing	pp 1–149 1
Pathophysiology - The Metabolic Response to Injury	1
<i>Acute Phase</i> Proteins:	1
Normal Wound Healing - The Three Phases	1
<i>Phase 2: Proliferation (Day3 to Week 3)</i>	2
<i>Phase 3: Remodelling (Weeks to Years)</i>	2
Fetal vs. Adult Wound Healing	2
Diagnosis & Management of the Acute Wound	3
Abnormal Scars	4
Chronic Non-Healing Wounds	4
Systemic Factors That Impair Healing	5
Surgical Site Infection (SSI) — CDC Wound Classification	5
Visual Patterns & Rapid Recognition	6
Pathophysiology — The "Rude Unhinging" of Life	7
2. Diagnosis — Classification and Recognition	7
SHOCK:Pathophysiology,Classification, andResuscitation	7
2.2 The Four Classes of Hemorrhagic Shock (ATLS) [HY]	8
2.3 Identifying Atypical Shock Patterns [HY]	8
2.4 Diagnostic Step-Ladder: Resuscitation Workup	9
3. Pathology — Shock Differentiators	10
4. Management — The Resuscitation Engine	11
5. Special Scenarios — Complications of Resuscitation	12
6. Visual Anchor Summary	12
SurgicalInfection:PathophysiologyandManagement	14
1. Pathophysiology - The Breakdown of Homeostasis	14
[HY] The Determinants of Outcome	14
2. Diagnosis - Classifying and Screening for Infection	15
CDC Operative Wound Classification [HY]	15
The Diagnostic Step-Ladder: Screening to Sepsis	16
3. Pathology - Tropical and Specific Mimics	17
4. Management - Sepsis and Antimicrobial Strategy	18
Antibiotic Stewardship and Prophylaxis	18
5. Special Scenarios - Catheter-Related Bloodstream Infection	18
6. Visual Anchor Summary	19
Fluids, Electrolytes & Acid-Base	21
Pathophysiology - The Metabolic Response to Injury	21
Body Composition & Fluid Compartments	21
Maintenance Fluid Therapy — The 4-2-1 Rule	22

Crystalloid Fluids — Comparison	23
Sodium Abnormalities	23
Potassium Disorders	24
<i>Management — in order:</i>	24
Hypernatremia ($\text{Na}^+ > 145 \text{ mEq/L}$)	24
Acid-Base Disorders	25
Hypokalemia ($\text{K}^+ < 3.5 \text{ mEq/L}$)	25
GI Losses & Acid-Base Consequences	26
Special Scenarios	26
Refeeding Syndrome	26

Nutrition and Metabolism in Surgical Patients..... 28

1. Pathophysiology — The Shift Into Autocannibalism	28
The Biphasic Response: Ebb vs Flow	28
2. Diagnosis — Working Out Who Needs Nutritional Support	29
4. Management — Routes and Requirements	30
3. Pathology — Starvation vs Trauma/Sepsis	30
Special Clinical Scenarios	31
Acute Pancreatitis	31
6. Visual Anchor Summary	32
Pathophysiology	33
The Metabolic Stress Response	33
Preoperative Evaluation & Risk Stratification	33

Preoperative Care..... 33

Stepwise Cardiac Evaluation	33
Scoring Systems	34
Hepatobiliary & Renal	35
Organ-Specific Risk Management	35
Pulmonary Optimisation	35
Risk factors:	35
Perioperative Medication Management	35
Continue vs. Hold	35
Intraoperative Safety — WHO Surgical Safety Checklist.....	36
Geriatric Patients & Frailty	36
Visual Patterns & Rapid Recognition	37
Pathophysiology - Why Any of This Matters	38
Preoperative Evaluation	38

Anaesthesia & Perioperative Care 38

Cardiac Risk Assessment — Step-by-Step	39
Airway Evaluation	40
ASA Physical Status Classification	40
Pharmacology — Induction & Maintenance	41
IV Induction Agents	41
Inhalational Agents	41
Neuromuscular Blockers (NMBs)	41
Intraoperative Management	42
WHO Surgical Safety Checklist	42
Postoperative Complications	43

💡 Heads up: Don't mix up Neurogenic Shock (a circulatory failure) with Spinal Shock (a temporary loss of spinal reflexes below the level of injury). Two different things that sound alike.

2.4 Diagnostic Step-Ladder: Resuscitation Workup

Goal is to restore perfusion before permanent end-organ damage (Multiple Organ Failure) sets in.

ABC Assessment
(Airway & Breathing first;
Shock management is "C")



Primary Survey
(Identify revealed vs concealed
hemorrhage sites)
→ bedside tests: FAST ultrasound,
Chest/Pelvis X-ray, or Diagnostic
Peritoneal Aspiration



Calculate Shock Index (SI) [HY]
→ $SI = HR / SBP$
→ $SI > 0.9$ suggests significant
bleeding / need for intervention
→ Modified SI (MSI) = HR / MAP is a
better predictor of mortality



Dynamic Fluid Challenge
(250–500 mL bolus over 5–10 mins)
→ categorize as: Responders,
Transient Responders (bleeding),
or Non-responders (exsanguinating)



Metabolic Perfusion Markers
(Arterial Blood Gas)
→ target: normalization of Base
Deficit (>6 mmol/L is severe)
and Lactate

3. Pathology

MAS Energy Devices and Risks


Comparative Clarity: MAS Energy Devices and Risks

Device/Condition	Mechanism	[HY] Key Clinical Differentiator
Monopolar Diathermy	Current through patient to return plate	Highest risk of capacitative coupling and insulation failure burns.
LigaSure [™]	Bipolar electrothermal energy + pressure	Can seal vessels up to 7 mm in diameter.
Harmonic Scalpel	Ultrasonic vibration (55.5 kHz)	Cuts and coagulates simultaneously with minimal lateral thermal injury.
CO ₂ Laser	Superficial absorption	Precise cutting but shallowest tissue penetration.
Nd:YAG Laser	Deep absorption	Deepest tissue penetration; used for bulky tumor destruction.



Heads up: Coupling injuries trip people up. Direct Coupling happens when the active electrode touches another metal instrument. Capacitative Coupling is an indirect injury where charge builds up between an insulated instrument and a conductive trocar, discharging into nearby tissue.


Comparative Clarity: Major C-Spine Fractures

Fracture	Image	Mechanism	Key Feature	Management
Jefferson		Axial Loading (Diving)	C1 burst fracture; lateral masses spread.	Stable (Transverse lig. intact): Hard collar. Unstable: Surgery.
Hangman		Hyperextension	C2 pars interarticular fracture; spondylolisthesis of C2 on C3.	Type I/II: Immobilization. Type III (facet dislocation): Posterior stabilization.
Odontoid		Variable	Fracture of the C2 dens (Type I, II, or III).	Type II (through base) is high risk for non-union; often needs surgery.

4. Management

Mandibular Fractures: Management

Open Reduction and Internal Fixation (ORIF) with titanium miniplates. Bilateral parasymphseal fractures require two miniplates on each side for stability.



Pediatric Fracture Patterns

Because children have more plastic bone and a substantial periosteum, they sustain unique injuries:

- Greenstick: Incomplete cortical disruption on the tension side.
- Torus/Buckle: Compression force causing the cortex to bulge.
- Plastic Deformation: Bowing of bone without a visible fracture line.

Comparative Clarity: Salter-Harris Classification (Growth Plate Injuries) [HY]

Type	Description	Prognosis/Management
I (S)	Slip: Straight across physis only.	Good; seldom affects growth.
II (A)	Above: Through physis and exiting metaphysis.	Most common type.
III (L)	Lower: Through physis and exiting epiphysis (Intra-articular).	Requires joint surface restoration.
IV (T)	Through: Across metaphysis, physis, and epiphysis.	High risk of growth disturbance; often needs surgery.
V (R)	Rammed: Crush injury of the physis.	Worst prognosis; often radiographically occult at first.

V. Comparative Clarity Tables

Suture Materials

Suture Material	Structure & Class	Tensile Strength Behavior in Tissue	Clinical Best Use
Polypropylene	Synthetic non-absorbable monofilament	Infinite tensile strength (resists enzymatic breakdown/hydrolysis)	Permanent tissue support; vascular anastomoses
Nylon	Synthetic non-absorbable monofilament	Gradual loss of strength over years via hydrolytic degradation	Skin closure; temporary fascial holding
Silk / Linen	Natural braided fiber	Rapid, progressive loss of strength via proteolytic degradation	Ligation of superficial vessels; temporary ties

Skin Grafts


Graft Type	Dermal Content	Metabolic Demands & Take Rate	Primary Cosmetic & Functional Outcomes
Split-Thickness (STSG)	Partial dermis	Low metabolic demands; higher take rate on poorly vascularized beds	Greater secondary contraction; suboptimal color/texture match
Full-Thickness (FTSG)	Complete dermis	High metabolic demands; requires highly vascularized bed	Minimal secondary contraction; superior color/texture match (face)

Diagnosis



The Diagnostic Step-Ladder: GERD to Malignancy

Rule out malignancy in anyone presenting with new-onset dysphagia.

Standard workup:



Heads up: In achalasia, the pH study can look abnormal too, but it's from fermentation of food residue sitting in a dilated oesophagus, giving "slow undulations" rather than the rapid reflux bursts you'd see in true GERD. Don't mix the two up.



Heads up: Ivor Lewis (two-phase — abdomen and right chest) is the standard for middle/lower third lesions. McKeown (three-phase — abdomen, right chest, neck) is the one to reach for with upper thoracic tumours, since you need that longitudinal clearance up in the neck.

Special Scenarios: Trauma and Perforation

Oesophageal perforation is a true surgical emergency.

Iatrogenic perforation [HY]: the most common cause (45–60%), usually during dilatation for achalasia or strictures

Boerhaave's Syndrome: spontaneous bursting (usually lower 1/3) after vomiting against a closed glottis



Management [HY]: Pittsburgh Severity Score aids prognosis; respiratory rate >30 with increasing O₂ needs carries high weightage

Mallory-Weiss Syndrome: partial-thickness mucosal tear at the cardia, presents with haematemesis




Visual Anchors: On-Sight Identification

"Bird's Beak": tapered distal narrowing, diagnostic of Achalasia

Heads up: SGLT-1 deficiency shows up as severe diarrhea even with breast milk, while GLUT-5 deficiency just causes bloating after eating fruit or honey.

Comparative Clarity: Regional Morphology

Feature	Jejunum	Ileum
Wall	Thicker	Thinner
Vasculature	Greater; long vasa recta	Less; short vasa recta
Arcades	Few	Many
Peyer Patches	Few	Most abundant



Small Bowel Obstruction: Mechanism

SBO is most commonly caused by postoperative adhesions.

Adhesions cause mechanical kinking → proximal dilation/distal collapse → bacterial translocation/ischemia.


Postoperative Ileus is different from mechanical SBO, it's functional. The inflammatory cascade starts with Mast Cells in the muscularis, which release NO and prostaglandins to paralyze smooth muscle.



The Appendix

1. Pathophysiology: From Obstruction to Perforation

The vermiform appendix is a midgut-derived, blind-ending muscular tube that arises from the cecum approximately 3 cm below the ileocecal valve. Previously considered vestigial, it's now recognized as a reservoir for "good" intestinal flora and a component of gut-associated lymphoid tissue (GALT).



Fundamentally, acute appendicitis is a disease of luminal obstruction. In adults, most commonly caused by a faecolith (appendicolith). Lymphoid hyperplasia is the primary trigger in children.


Obstruction leads to a predictable cascade:

Ongoing mucus secretion increases intraluminal pressure
Impairs venous drainage, causing mucosal ischemia and bacterial translocation
If it continues, full-thickness ischemia results in gangrene and perforation, releasing a polymicrobial inoculum (most commonly *E. coli* and *Bacteroides fragilis*) into the peritoneal cavity

This correlates with the hallmark clinical triad: early vague periumbilical pain (visceral T10-L1 stimulation), followed by anorexia/vomiting, and culminating in localized right lower quadrant (RLQ) pain (somatoparietal irritation).

2. The Diagnostic Step-Ladder: From Scoring to Imaging

The diagnosis remains essentially clinical, but a systematic workup reduces the negative appendectomy rate, which is traditionally higher in women of childbearing age (up to 25-30%).



2. Diagnostic Approach

The Diagnostic Step-Ladder

For any suspicious breast symptom, the Triple Assessment is the mandatory workup path. When all three components are concordant, the Positive Predictive Value exceeds 99.9%.

Step-Ladder Flowchart:

Clinical History & Physical Exam
(Inspection → Palpation)



Radiological Imaging
(Age <35: Ultrasound
Age >35: Mammography)



Tissue Sampling
(FNAC for cytology, or
Core Biopsy for architecture)





Multidisciplinary Team (MDT)
Review for Management

Heads up: While Ultrasound is excellent for distinguishing cysts from solid lesions in young women with dense breasts, Mammography remains the gold standard for population-based screening in women over 40–50 years.

Conn's Syndrome: Workup and Localization

Workup: PAC/PRA Ratio >30 is highly suggestive.

Localization Step-Ladder:



Heads up: CT is the first imaging step, but AVS is required in patients >40 years because incidental non-functioning nodules are common in this age group and can lead to false-positive CT results.

Cushing's Syndrome (Hypercortisolism): Diagnosis

Screening: 24h urinary free cortisol (at least twice), late-night salivary cortisol, or 1mg overnight DST.

Etiology: Measure plasma ACTH. Low ACTH (<5 pg/mL) = Adrenal source. High ACTH = Pituitary or Ectopic source.

Pheochromocytoma: Biochemical Diagnosis

Plasma free metanephrines (High sensitivity) or 24h urinary metanephrines/catecholamines (High specificity).

Heads up: Never biopsy a suspected adrenal mass until pheochromocytoma has been biochemically excluded. Biopsy can trigger a fatal hypertensive crisis. Also, never give beta-blockers first, it causes unopposed alpha-stimulation and severe hypertension.

3. Pathology

Comparative Clarity: Varieties of Hyperparathyroidism

Feature	Primary (PHPT)	Secondary (SHPT)	Tertiary (THPT)
Mechanism	Autonomous adenoma	Response to low Ca (Renal Failure)	Autonomous after Renal Transplant
Calcium	High	Low or Normal	High
PTH	High	Very High	High
Key Sign	Kidney Stones	Renal Osteodystrophy	Persistent Hypercalcemia post-transplant

Bone metastasis follows the "Seed and Soil" theory — tumor cells (the seed) get recruited to the bone marrow (the soil) via the CXCL12/CXCR4 chemokine pathway.

Clinical Etiology and Risk Factors

Environmental and modifiable factors drive an estimated 60–90% of cancer cases.

Tobacco smoking is the biggest driver. A "Current Smoker" is defined as someone who has smoked at least 100 cigarettes in their lifetime and currently smokes ≥ 1 cigarette per day. Smoking induces a higher tumor mutational burden (TMB), which increases mortality but can actually make some tumors (like lung cancer) more responsive to PD-1/PD-L1 immunotherapy, since higher TMB means more neoantigen expression.


Comparative Clarity: Major Modifiable Risk Factors [HY]

Risk Factor	Most Strongly Associated Cancer	Key Exam Fact
Obesity	Endometrial Cancer	RR is far higher than other obesity-related cancers.
Alcohol	Mouth, Pharynx, Larynx, Esophagus, Breast	Any amount of alcohol increases risk; higher intake adds stomach/liver risk.
Smoking	Lung, Oropharynx, Bladder	Multiplexes risk of second primary cancers if continued during CT/RT.
UV Radiation	Melanoma, BCC, SCC	Melanoma has the highest mutational burden of any studied malignancy.

Comparative Clarity: Renal Cortical Subtypes [HY]

Feature	Clear Cell RCC (70-80%)	Papillary RCC (10-15%)	Chromophobe RCC (3-5%)	Oncocytoma (Benign)
Genetic Driver	VHL (3p deletion)	MET (Type 1); FH (Type 2)	Loss of Y, 1, 2, 6, 10	Benign
Gross Appearance	Golden-yellow (lipid-rich)	Tan-brown, friable	Pale, beige, translucent	Mahogany-brown
Key Clinical Factor	Most common; respond to immunotherapy	Multifocal; poor prognosis if multifocal	Excellent prognosis	Central stellate scar
IHC / Special	CA-IX overexpression	-	CK7 Positive	CK7 Negative


Bladder/Urothelial Carcinoma Workup



Comparative Clarity: Prostate Biopsy Interpretations [HY]

Finding	Gleason Score	Grade Group (GG)	Management Path
Well-formed glands	3 + 3 = 6	GG 1	Active Surveillance
Differentiated/Aggressive mix	3 + 4 = 7	GG 2	Surgery or RT + Short ADT
Aggressive/Differentiated mix	4 + 3 = 7	GG 3	Worse prognosis than GG2
Poorly formed/Cribriform	8	GG 4	Multimodal: RT + Long ADT
Sheets/Necrosis	9/10/2026	GG 5	Multimodal: RT + Long ADT

Testicular GCT Workup



Heads up: Transscrotal biopsy or orchiectomy is strictly contraindicated, it risks tumor cell seeding into inguinal and pelvic lymphatics.

Comparative Clarity: Serum Marker Profiles [HY]

Marker	Seminoma	Non-Seminomatous (NSGCT)	Half-Life
AFP	Always Normal	Elevated (Yolk Sac / Embryonal)	5-7 Days
hCG	Elevated in 10%	Elevated (Choriocarcinoma / Embryonal)	2-3 Days
LDH	Marker for bulk, not prognosis	Prognostic factor	Variable


3. Management RCC

Partial Nephrectomy (PN): standard of care for T1 masses (<7 cm) and "absolute" indications (solitary kidney, bilateral tumors, CKD, or familial RCC).

Radical Nephrectomy (RN): reserved for high-complexity tumors where PN isn't feasible, provided the contralateral kidney is normal and baseline eGFR is >45. Dissection is performed external to Gerota's fascia, and the adrenal is spared unless directly invaded.

Advanced/Metastatic: Tyrosine Kinase Inhibitors (TKIs) like sunitinib/pazopanib (VEGF inhibition) are the backbone, often combined with Immune Checkpoint Inhibitors (ICIs).

Diagnostic step-ladder:



Management hierarchy:

Conservative: 90% of stones <5 mm pass spontaneously. Medical Expulsive Therapy (Tamsulosin) relaxes ureteral smooth muscle.

Surgical intervention: indicated for persistent pain, infection, or size >6-10 mm.


ESWL: best for renal stones <2 cm. [HY] Complication: Steinstrasse (stone street), ureteral obstruction by fragments.

PCNL: preferred for larger stones (>2 cm) or staghorn calculi.

Emergency drainage: if sepsis is present with obstruction, urgent Percutaneous Nephrostomy (PCN) is safer than a stent.

Heads up: Non-contrast CT is the most sensitive test overall, but pure uric acid stones are radiolucent on X-ray, they only show up on CT. Uric acid stones are the unique ones you can actually dissolve, by raising urinary pH to 6.5 with potassium citrate.

Clinical clarity: trauma classification [HY]

Injury Type	Image	Mechanism	Diagnostic Finding	Management
Intraperitoneal Bladder Rupture		Blow to a full bladder	Contrast outlining bowel loops/peritoneal cavity	Emergency Surgical Repair
Extraperitoneal Bladder Rupture		Pelvic fracture	Contrast limited to perivesical space	Conservative (Catheter drainage for 10-14 days)
Bulbar Urethral Injury		Straddle/Fall-astride	Butterfly hematoma in perineum	Suprapubic Catheter (SPC) + Delayed Repair
Membranous Urethral Injury		Pelvic fracture	High-riding prostate on DRE; RUG showing extravasation	Suprapubic Catheter (SPC); avoid Foley

Heads up: Extraperitoneal ruptures are usually managed conservatively, but surgical repair is mandatory if there are bone spicules in the bladder or a concomitant rectal injury.


6. Oncology: Bladder and Urethral Carcinoma

Bladder cancer is the 5th most common adult malignancy, with painless gross hematuria being the most common presentation.

Non-Muscle Invasive (NMIBC: Ta, T1, Tis): limited to mucosa or lamina propria. Management: TURBT (Gold Standard) + Intravesical therapy (BCG or Mitomycin C).

Diagnostic step-ladder for LUTS/BPH:


The goal is to differentiate benign enlargement from malignancy and evaluate for "high-pressure" complications.



Heads up: Pressure-flow urodynamic studies are NOT essential for every patient. Save them for complex cases where symptoms don't match the flow rate, or a neurological cause is suspected.

1. Pathophysiology: The "Rigid Box" and Secondary Insult

Neuro trauma is fundamentally a battle against the physics of the Monro-Kellie doctrine. The cranium is a "rigid box" with a fixed internal volume occupied by brain parenchyma, blood, and cerebrospinal fluid (CSF). Because these components are nearly incompressible, an increase in any one, whether from a hematoma, brain swelling, or hydrocephalus, must be offset by a decrease in the others.




Pathology is divided into:

Primary Injury: the immediate mechanical damage to neurons and vessels at impact.

Secondary Injury: the true target of neurosurgical intensive care. A cascade of damage caused by hypoxia ($\text{PaO}_2 < 60 \text{ mm Hg}$) and hypotension ($\text{SBP} < 90 \text{ mm Hg}$), which set off inflammation, calcium influx, and cell death.

When compensatory displacement of CSF and venous blood is exhausted (the point of decompensation), intracranial pressure (ICP) rises exponentially, leading to rostrocaudal decay and the classic hallmark clinical triad (Cushing's Triad): hypertension, bradycardia, and irregular respirations, signaling imminent brainstem herniation.




2. The Diagnostic Step-Ladder


The workup for neuro trauma follows a rigid hierarchy to prioritize "life over limb" and prevent secondary injury.

I. Principles of Stereotaxy and Radiosurgery Pathophysiology

Functional neurosurgery runs on the Monro-Kellie doctrine too — cranium as a rigid vault with fixed volumes of brain, blood, and CSF. Functional disorders come from hyperfunction or deranged activity in specific neural circuits. To hit these deep, invisible targets, surgeons use stereotaxis — localizing a target in 3D space with a calculated coordinate system (x, y, z).



Diagnostic step-ladder for stereotactic procedures:




CNS Infections

1. Pathophysiology

Intracranial Dynamics and the "Rigid Box"

CNS infections are surgical emergencies not just because of pathogen virulence, but because of the Monro-Kellie doctrine. The adult cranium is a "rigid box" with a fixed internal volume. Any infectious process, whether the diffuse inflammatory swelling of meningitis, the localized pus collection of a brain abscess, or the vasogenic edema of cerebritis, adds volume to this box.

Once the brain's compensatory reserves (displacement of CSF and venous blood) are exhausted, intracranial pressure (ICP) rises exponentially. This produces the hallmark clinical triad of intracranial hypertension: hypertension, bradycardia, and irregular respirations (Cushing's Triad). Left untreated, this pressure gradient drives rostrocaudal decay, where brain tissue is forced through narrow internal openings (herniation), leading to brainstem compression, midbrain infarction (Duret hemorrhages), and death.



2. Diagnosis

The Diagnostic Step-Ladder

The workup for a suspected CNS infection must prioritize identifying life-threatening mass effect before proceeding to definitive pathogen identification.

Clinical Suspicion
(Fever, headache, and "meningism")



Emergent Non-contrast Head CT
(mandatory before lumbar puncture
to rule out mass effect or
hydrocephalus and prevent
terminal "coning"/downward
herniation)

Special Scenarios & Clinical Pearls

Heads up: Clinical evaluation - colour, capillary refill, warmth - is the gold standard for monitoring free flaps post-op. The most critical window is the first 48 hours.

Heads up: The "delay" phenomenon increases flap survival by enlarging existing arteries along the flap axis via cellular hypertrophy. Maximal effect kicks in between 48 and 72 hours.

Marjolin's Ulcer - rare, highly aggressive squamous cell carcinoma arising in chronic wounds or scars (burns, osteomyelitis). High propensity for metastasis.



Ideal dressing - must provide a moist environment, enable gas exchange, and protect against infection.

Photography standards - same photographer, camera, and lighting for pre- and post-op photos.

Visual Anchors

Langer's Lines / RSTLs - the "natural wrinkles" of the face. Incisions parallel to these are nearly invisible; incisions across them create wide, unfavourable scars.




Classification by Depth

Depth	Layer Involved	Clinical Appearance	Sensation	Healing / Management
Superficial (1st degree)	Epidermis only	Erythematous, dry, no	Painful	Spontaneous healing in 3–6
Superficial Partial-	Epidermis + Papillary	Bright red, moist,	Exquisitely painful; hair	7–14 days; minimal
Deep Partial-Thickness	Epidermis + Reticular	Pale, mottled,	Reduced sensation;	>21 days; often
Full-Thickness	Entire dermis +	Leathery eschar,	Anaesthetic (painless)	Requires excision and

Heads up: Laser Doppler and AI tools are emerging, but the gold standard for assessing burn depth is still serial clinical judgement by an experienced practitioner.


Initial Assessment & TBSA Calculation



Airway assessment - evaluate the pharynx for oedema or particulate immediately upon extraction from a closed-space fire.

4. Craniosynostosis: Virchow's Law and Skull Morphology

Craniosynostosis = premature fusion of one or more cranial sutures.
 Mechanism (Virchow's Law): growth is restricted perpendicular to the fused suture, with compensatory growth parallel to it.



The diagnostic step-ladder:

[Physical Exam]
 (98% accurate — look for palpable ridges/dysmorphic shape)




[3D Computed Tomography]
 (Gold standard for definitive diagnosis)



[Intracranial Pressure (ICP) Screening]
 (Required for all, especially multisuture cases)

Suture Fused	Deformity Name	Physical Presentation
Sagittal (Most Common)	Scaphocephaly	Long, narrow skull; frontal/occipital bossing.
Metopic	Trigenocephaly	Triangular (keel-shaped) forehead; hypotelorism.
Bicoronal	Brachycephaly	Short, wide skull; flattening of occiput.
Unilateral Coronal	Anterior Plagiocephaly	Orbital asymmetry and unilateral frontal flattening.
Multiple Sutures	Cloverleaf (Trilobar)	Severe deformity; surgical emergency due to hydrocephalus/ICP.



Metabolic Acidosis: What It Signals

Usually signals inadequate tissue perfusion (e.g., shock, NEC, or volvulus).

3. Pathology

Comparative Clarity: TPN vs. Enteral Feeding

Feature	Total Parenteral Nutrition (TPN)	Enteral Feeding
Primary Risk	PN-Associated Liver Disease (PNALD) / Cholestasis	Malabsorption / Stressed gut
Metabolic Sign	Rising Bile Acids → Rising Direct Bilirubin	Best physiological stimulus for bile flow
Cholestasis Prevention	Use Omega-3 fatty acid emulsions (Omegaven)	Early restoration of enteral feeds is most effective


First Branchial Cleft Anomalies

Located near the external auditory canal, front or back of the ear, or the submandibular region near the mandible angle. Tract usually runs through or deep to the parotid gland, wrapping around or passing between branches of the facial nerve (CN VII) before ending in the external auditory canal. Complete excision needs careful preservation of CN VII.

Second Branchial Cleft Anomalies (most common, 70-95%)

Classic presentation: lateral neck mass or draining sinus at the lower anterior border of the SCM (junction of upper half and lower third).

Tract path (high yield): penetrates the platysma, ascends along the carotid sheath to the hyoid bone, turns medially, and runs between the internal and external carotid arteries. Then passes close to the hypoglossal nerve (CN XII) and glossopharyngeal nerve (CN IX) (behind the posterior belly of digastric and stylohyoid) before ending in the tonsillar fossa of the lateral pharyngeal wall.



External Ostium (Lower 1/3 anterior SCM)



Penetrates Platysma



Ascends along Carotid Sheath



Reaches Hyoid Bone and turns medially



Passes between Internal & External Carotids (Carotid Bifurcation)



Crosses adjacent to CN XII (Hypoglossal) and CN IX (Glossopharyngeal)



Internal Ostium (Tonsillar Fossa / Lateral Pharyngeal Wall)

Surgical strategy: pass a fine lacrimal probe or inject methylene blue into the tract to help with dissection. Removing this long, winding tract completely needs multiple small, staggered transverse incisions (stepladder technique) along the anterior SCM border. Gives better exposure, less tissue trauma, and better cosmetic result than one long oblique incision.



Classification [HY]: Type I (75%) consists of large cysts (2–10 cm) and has a good prognosis; Type III is microcystic/solid and carries a poor prognosis due to lung hypoplasia and hydrops.

Vascularity: Characteristically supplied by the pulmonary artery and drains via pulmonary veins.


Pulmonary Sequestration (BPS) Non-functioning lung tissue with no connection to the tracheobronchial tree.

Classic Marker [HY]: Derives its blood supply from an aberrant systemic artery (usually the aorta).

Comparative Clarity: Congenital Lung Lesions

Condition	Image	Key Differentiator	Vascular Supply
CPAM		Multiloculated cysts; communicates with airway	Pulmonary Artery
Sequestration		Non-functional tissue; no airway connection	Aberrant Systemic Artery (Aorta)
CLE		Overdistension of a single lobe	Normal Pulmonary Vessels
Bronchogenic Cyst		Ciliated columnar lining; typically carinal/hilar	Normal Bronchial Vessels

Staged palliation hierarchy [HY]:



Obstructive & Miscellaneous Defects


Coarctation of the Aorta (COA): narrowing distal to the left subclavian origin
High-yield exam: radiofemoral delay and significant BP discrepancy between arms and legs

Anomalous Left Coronary Artery from the Pulmonary Artery (ALCAPA):
Bland-White-Garland syndrome. Most common cause of myocardial ischemia in infants, treated by surgical reimplantation into the aorta

Eisenmenger's Syndrome: late reversal of L-to-R shunt to R-to-L from irreversible PVR. Surgical repair is contraindicated at this stage

Visual Anchors

Tetralogy of Fallot: "boot-shaped heart" (coeur en sabot) on CXR from RVH and concave pulmonary segment




TGA: "egg-on-a-side" (egg-on-string) silhouette with narrow mediastinum



Symptomatic: pressure on nearby structures gives hoarseness (recurrent laryngeal nerve), dysphagia (esophagus), stridor (bronchus)
Most common location [HY]: aortic root and ascending aorta

Diagnostic path



Heads up: CXR can show a widened mediastinum, but a normal CXR happens in up to 16% of aortic dissections, so it can't rule anything out. CTA is your standard for stable patients, but if the patient's unstable, TEE is the most sensitive bedside test.

Management

Surveillance — small, asymptomatic aneurysms

Medical — aggressive BP control (beta-blockers)

Surgical thresholds [HY]:

Sporadic ascending TAA: ≥ 5.5 cm

Descending TAA: ≥ 6.0 cm

Marfan/syndromic: lower threshold (4.5–5.0 cm), or if expanding ≥ 0.5 cm/year

Bicuspid aortic valve (BAV): intrinsic aortopathy risk, repair if > 4.5 cm and having other cardiac surgery anyway

Heads up: In the Fontaine classification, Stage IV means ischemic ulcers only. Venous ulcers never make it into this system, don't mix them up on an exam.

5. Management

Initial management — Intermittent Claudication

Goal is symptom relief plus cardiovascular risk reduction.

First-line [HY]: Supervised Exercise Therapy (30-45 mins, 3x/week for 12 weeks) plus aggressive risk factor modification (smoking cessation, statins, BP control).

Pharmacotherapy:

Cilostazol (PDE-III inhibitor): the only FDA-approved drug that increases walking distance.

Aspirin/Clopidogrel: reduce cardiovascular events (MI/stroke) but don't improve walking distance.

Heads up: Cilostazol is contraindicated in Heart Failure, it's linked to increased mortality in that group [HY]. Don't reach for it just because it's the walking-distance drug.

Revascularization Strategy

Indicated for disabling claudication unresponsive to medical therapy, or CLI/limb salvage.

Iliac (aortoiliac): endovascular angioplasty and stenting often preferred.

Femoro-popliteal (SFA):

Short lesions (<10-15 cm): endovascular therapy is first-line.

Long-segment occlusions (>15 cm) [HY]: femoropopliteal bypass with autologous vein graft (great saphenous vein) is the gold standard for durability and patency.

ALI management: urgent surgical thromboembolectomy using a Fogarty balloon catheter for embolic occlusions with viable runoff.